



BROKEN BONES TREE

General Details		
Reference	https://biophilia.lk/database/plant/broken-bones-tree	
Scientific Name	Oroxylum indicum	
Sinhala Name	????? (THOTILA)	
Tamil Name	???????????? ???? (PUTAPUSPAM MARAM)	
Other Common Names	Midnight Horror, Oroxylum, Indian Trumpet Flower, Broken Bones, Indian Caper, Tree of Damocles	
Synonyms		
Plant Family	Bignoniaceae	
Plant Group	Angiosperms (Enclosed Seed)	
Plant Types	Medium Tree	
Morphologies / Crown Architecture	Open Canopy	

Categories	Canopy plants Dry Zone Plants Edible Plants Flowering Plants Forest Trees Fruiting Plants Herbal Plants Intermediate Zone Plants Ornamental Plants Fast Growing Plants Sun Loving Plants Trees Wild Plants
Native Regions	Indian Subcontinent
Distributions	E. Asia - India, Sri Lanka, Thailand, Malaysia, Indonesia, Philippines, New Guinea, northern Australia, New Caledonia, Vanuatu.
Habitat	Lowland woodlands, clearings and waysides. Tropical and subtropical low altitude open forests, roadsides, slopes
Elevation	Above sea level up to 1,000 m
Average Height	10.00
Maximum Height	20.00
Root Systems	Tap Root

Propagations	By Seeds By Cutting (Stem)		
Indoor	N/A		
Medicinal Use	Yes		
Medicinal Value	Stem Bark and Roots for Diarrhoea, Dysentery, Rheumatism, Otorrhoea and Piles in Sinhala Ayurveda		
Special Features	Young leaves and flowers - raw or cooked Eaten uncooked as a side dish with rice Fruit - cooked. Young fruits are cut into pieces, boiled and eaten with rice The flowers, which open of a night time, have a foetid smell that is attractive to bats		
Foliage	Foliage		
Leaf / Habit	Evergreen		
Leaf Shape	Obtuse shape		
Arrangements	Simple		
Leaf Arrangement	Odd Pinnate Leaf Arrangement		
Flowering			
Is Flowering ?	Yes		
Blossoms	Large, fleshy, and foul smelling with 2.5-cm-long pedicle. Calyx and corolla are about 2.5 cm and 6.5 cm in size, respectively. Stamens are slightly exerted and base of filaments is wooly.		
Blooming Season	July; August		
Fruiting			
Bears Fruits ?	Yes		

Fruit	Follicle, 30–90 cm long and 5–10 cm broad, strap/sword shaped, compressed, and two-valves. Fruiting Time: From December to March Seed: Flattened and winged.	
Maintenance		
Level	Medium	
Soil Types	Sandy Soil Loam Soil Well Drained Soil	
Water	Medium/Moderate	
Sunlight	Full	
Humidity	Average	
Edibility		
Is Edible ?	N/A	
Legal Status		
Flora Fauna Protection Ordinance 2009 Amendment, Sri Lanka	Not Evaluated	